

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):


- BLACK BORDERS
- ✓ • TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**


1/7

FIG. 1A




R = $-\text{C}(\text{O})\text{NH}_2$ (1a);
 $-\text{C}(\text{O})\text{NHCH}_3$ (1b);
 $-\text{C}(\text{O})\text{N}(\text{C}_2\text{H}_5)_2$ (1c);
 $-\text{C}(\text{S})\text{NH}_2$ (1d);
 $-\text{C}(\text{O})\text{CH}_3$ (1e);
 $-\text{C}(\text{O})\text{CCH}_3$ (1f);
 $-\text{CN}$ (1g);
 $-\text{CH}_3$ (1h);
 $-\text{H}$ (1i)

FIG. 1B



2/7

FIG. 1C



substrate	R	relative rate ^A	turnover/h ^B
1a		1.0	8
1b		0.9	8
1c		0.0	0
1d		1.3	11
1e		1.1	9
1f		1.3	11
1g	-CN	0.9	8
1h	-CH ₃	0.0	0
1i	-H	0.0	0






FIG. 4

4/7







FIG. 5

FIG. 6









5/7

Mimic NADH Models in Biocatalysis
 Tandem Cofactor Regeneration, Enzyme Recognition and
 Chiral Synthesis of Alcohols



6/7

Enzymatic Reductions of Ketones with NAD⁺ Models:
 Turnover Frequencies and Enantiomeric Excess^{a,b}


substrate	product	% yield	TOF(d ⁻¹)	ee (%S)
		90(91)	30(31)	93(93)
		55(59)	18(19)	>99(99)
		5(5)	4(4)	>96(96)
		41(59)	14(20)	85(85) ^c

^aThe results from NAD⁺ were given in parenthesis. ^bThe enantiomeric excess was determined by GLC with a modified β -cyclodextrin capillary column. ^cBased on derivatization with an optically pure isocyanate.

FIG. 7

7/7

Biomimetic Reduction of Norcamphor



^a The result for NAD⁺ was given in parenthesis.

FIG. 8




FIG. 9